

Vol 2 / Issue 1 / January-March 2011

Chronicles of Young Scientists



The use of antibiotics and disinfectants in the freshwater prawn (*Macrobrachium rosenbergii*) farms of north and south 24 Parganas districts of West Bengal

Abstract

The freshwater prawn (*Macrobrachium rosenbergii*) farms and hatcheries of north and south 24 Parganas districts in West Bengal were surveyed for the details of antibiotics and disinfectants used for controlling the prawn diseases and pathogens. During the survey, 4antibiotics and four 4disinfectants were found to be used commonly by the farmers during a 12-month period of study. The chemicals used in the culture systems as antibiotics or disinfectants for the culture water, tanks, other utensils, and others have serious health hazards for both the cultured prawns and human beings. The study also indicates no herbal medicines, and probiotics are being used in prawn farms.

Key words:

Antibiotics, disinfectants, freshwater prawn farm, *Macrobrachium rosenbergii*

Introduction

Polyculture in freshwater fish–prawn farms, prawn farms, and rice–prawn farming are the tradition of West Bengal, especially in the delta regions of north and south 24 Parganas districts. With the growing demand in domestic markets as well as in foreign markets, the culture of freshwater prawn has proved its economic importance and become burgeoning business in West Bengal. The giant freshwater prawn, *Macrobrachium rosenbergii*, De Man,^[1] locally known as “*Golda Chingri*” (Bengali), plays an important role in the economy of West Bengal by earning valuable foreign exchange, generating employment, improving the socioeconomic conditions of the backward people and raising protein-based food production besides the culture of fishes, paddy, and other organisms. Nevertheless, the biggest problem faced by the farmers every year in this sector due to the viral, bacterial, or parasitical diseases. The farmers do not give any treatment for viral diseases that attack the stock, as these are incurable. The bacterial diseases are treated by different antibiotics to avoid the

outbreak of deadly diseases. The quality of the water and soil is always maintained by using disinfectants.

The present study deals with the antibiotics and chemicals that are used in the freshwater prawn farms and hatcheries in two districts (north and south 24 Parganas) of West Bengal for the control of disease outbreaks.

The prawn farming industry has been subjected to criticism from an environmental point of view. Concern has been expressed regarding the use of chemicals in farms and its potential impacts on the environment and human health.^[2-4]

Materials and Methods

The survey was conducted during January 2009 to February 2010 in different fish–prawn farms based on freshwater

Monjit Paul, Mukti Chanda, Joydev Maity¹,
Supriya Sen Gupta, Bidhan Chandra Patra², Gadadhar Dash³

Department of Industrial Fish and Fisheries, Asutosh College, Kolkata, ¹Departments of Aquaculture Management and Technology and ²Zoology, Vidyasagar University, Midnapore, ³Department of Fish Pathology, Faculty of Fishery Science, West Bengal University of Animal and Fishery Sciences, Kolkata, West Bengal, India

Address for correspondence:

Dr. Monjit Paul,
P - 227/1, Upen Banerjee Road, Kolkata - 700 060,
Parnashreepally, Behala, West Bengal, India.
E-mail: monjit.paul@gmail.com

Access this article online	
Website: http://www.cyonline.org	Quick Response Code 
DOI: 10.4103/2229-5186.79352	

wetlands (bheries) in 2 districts (north and south 24 Parganas) of West Bengal in India. About 50 farms, including wetland culture systems were surveyed and the information related to the use of different chemicals, drugs, antibiotics, herbal medicines, and probiotics used in the farms or wetland culture systems were collected by the basis of personal interview with different farmers or members of the cooperative societies culturing the prawns by predefined questionnaire.

Result

A variety of drugs are being used predominantly in different farms of the districts to reduce the attack of the different pathogens, to improve the survival rate and to increase the intensity of feeding leads to better growth rate. The common antibiotics oxytetracycline, erythromycin, chloramphenicol, nitrofurans were found to be used to control the bacterial pathogens successfully. Most of the farms use chloramphenicol (40%) followed by oxytetracycline (23%) and erythromycin (20%).

Besides these antibiotics, the farms use disinfectants, such as formalin, malachite green, potassium permanganate, and methylene blue (MB), to disinfect the water and for parasitical diseases caused by petrichous protozoan ciliates, and some bacterial pathogens [Table 1]. No fish farms of two districts were found to use any herbal medicines and probiotics to treat the prawns or their seeds.

Discussion

The chemicals that being used in the prawn farms and

the hatcheries are effective mostly prior to the infection, or at the preliminary stages of the infection. Mostly the farmers used prophylactic treatment measures based on eye observations of clinical signs. Food and Drug Administration (FDA)^[5] legalized 5 drugs/chemicals in US aquaculture, such as oxytetracycline, sulfamerazine, ormetoprim, formalin, tricaine methanesulfonate. FDA^[6,7] legalized the following drugs for using as animal drugs: Finquel (MS-222), Tricaine-S (MS-222), Formalin-F, Paracide-F, Parasite-S, Romet 30, Sulfamerazine in Fish Grade, Terramycin-200, Chorulon, 35% PEROX-AID, Aquaflor, Aquaflor-CA1, TERRAMYCIN-343, Oxytetracycline HCl Soluble Powder-343, OxyMarine, TETROXY Aquatic, and Terramycin-200 for Fish.

Theoretically, chemicals other than antibiotics that are added to the shrimp ponds, or byproducts from the applied substances, that have a bioaccumulation potential, could be found as residues in the shrimps. However, little attention has been paid to the risk of residues other than antibiotics in farmed shrimps, and no data from such investigations have been found, Gräslund and Bengtsson.^[3]

The antibiotics chloramphenicol (mostly used in the farms) and nitrofurans are banned worldwide for use in the production of foods because of their serious side effects. Chloramphenicol may cause fatal aplastic anemia and nitrofurans are classified as carcinogens.^[3,8,9]

Aftab Uddin *et al.*^[9] reported to control the uses of antibiotics in prawn farms due to the risk of development of resistant bacteria. When the bacteria have acquired resistance, it is

Table 1: Antibiotics and disinfectants used in the prawn farms of north and south 24 Parganas districts of West Bengal during the study period (January 2009 to February 2010)

Antibiotics/disinfectants	Farms visited (%)	Dosage (ppm)	Diseases	Pathogens
Chloramphenicol	40	10	Filamentous bacteria Luminous bacteria	<i>Leucothrix mucor</i> <i>Vibrio harvey</i> <i>Vibrio splendidus</i>
Oxytetracycline	23	10 3-5	Vibrio infections Bacterial shell diseases	<i>Vibrio parahaemolyticus</i> <i>Vibrio anguillarum</i> <i>Vibrio</i> spp., <i>Aeromonas</i> spp. <i>Pseudomonas</i> spp.
Erythromycin	20	2-4	Luminous bacteria	<i>Vibrio harvey</i> <i>Vibrio splendidus</i>
Nitrofurans	10	1-2	Black gill disease	
Formalin	81	10-25	Protozoan ciliates	<i>Zoothamnium</i> spp., <i>Epistylis</i> spp., <i>Vorticella</i> spp., <i>Acineta</i> spp.,
Malachite green	34	0.075 0.0075	Luminous bacteria Shell diseases Protozoan ciliates	<i>Vibrio harvey</i> <i>Vibrio splendidus</i> <i>Vibrio</i> spp., <i>Aeromonas</i> spp., <i>Pseudomonas</i> spp.
Formalin + malachite green	22	0.01 + 0.01	Bacterial diseases, protozoan diseases	<i>Zoothamnium</i> spp., <i>Epistylis</i> spp., <i>Vorticella</i> spp., <i>Acineta</i> spp.,
Potassium permanganate	14	3-5	Luminous bacteria, Shell diseases	
Methylene blue	35	8-10	Protozoan ciliates	<i>Zoothamnium</i> spp., <i>Epistylis</i> spp., <i>Vorticella</i> spp., <i>Acineta</i> spp.,
Herbal medicines and probiotics	0	-	-	-

impossible to get rid of them with antibiotics that caused the resistance.

Frappaolo *et al.*^[10] reported that the use of antibiotics to control bacteria population and maintain a healthy environment for prawn culture becomes popular. A wide range of antibiotics is now being used to treat bacterial diseases and to control bacterial population in the hatcheries and prawn farms. The use of antibiotics for treatment arise various antibiotic-resistant bacteria. The resistance was transfer to pathogenic bacteria, and led to reduce efficacy of antibiotic treatment for disease caused by the resistant pathogens.

The antibiotics when used in the prawn farms create health hazards when exposed on the skin or inhaled.^[9] Same problem was observed in the fish farms of 2 districts, where the workers suffered skin irritation and respiration problems due to the use of such chemicals.

Malachite green is a respiratory poison reported by Dieberg and Kiattisimkul.^[11] Malachite green is not very soluble in water, and it binds to sediments reported by Bergheim and Asgard.^[12] It accumulates in biota but mostly in simple organisms. In fish, malachite green can be found in all organs in great quantities, including kidney.^[13] Residues of 2400 µg/g have been found in fish, and they are persistent.^[14]

In water, potassium permanganate is quickly transformed into nontoxic manganese dioxide, which precipitates out. It is toxic for phytoplankton reported by Reardon and Harrell,^[15] and Gräslund and Bengtsson.^[3]

MB at low concentrations in the body, exist in equilibrium and form a reversible oxidation–reduction system, which is the basis for MB's function as an electron donor/acceptor and free radical scavenger. As such, MB has the potential to affect a wide variety of physiologic reactions. Its effect on red blood cells is to reduce the heme from methemoglobin to hemoglobin. Paradoxically, at higher doses, this equilibrium is destroyed and an excess of methemoglobin is produced as reported by Curry.^[16] The formation of hydrogen peroxide as a byproduct can lead to oxidation of the red-cell membranes, denaturation of hemoglobin, hemolytic anemia, and Heinz body formation.^[17,18] Recent research has focused on the vasoactive properties of MB. In response to agents, such as acetylcholine and bradykinin, the vascular endothelium produces nitric oxide (NO) through the action of NO synthase. NO activates soluble guanylate cyclase, which raises the levels of cyclic guanosine monophosphate (cGMP), opening calcium-sensitive potassium channels and producing membrane hyperpolarization, smooth muscle relaxation, and vasodilatation.^[19-22]

Formalin is toxic to aquatic life at low concentrations, with

96 h LC 50 of 1–1000 µL/L.^[8] Some fish are sensitive to it, so a bioassay is recommended before use as reported by Noga.^[23] Inhibitions of growth and mortality have been reported for phytoplankton and macrophytes (aquarium plants). Its algicidal property can further reduce oxygen. Toxicity is more important in low pH water and at a high temperature. When stored at room temperature, formalin can develop a white precipitate of paraformaldehyde, which is more toxic than pure formalin as reported by Douet *et al.*^[24]

Disinfectants are drained in the local water bodies lead to pollution. The problems can be eliminated by implementing laws to control the hazardous drugs. The farms may use procedures to avoid the introduction of diseases in prawns by regular water quality management, probiotics, or other herbal medicines with fewer side effects. If the farms are infected with pathogens leading to diseases, the prawns must be treated with approved chemicals.

References

1. De Man JG. On some species of the genus *Palaemon* Fabar. With descriptions of two new forms. Notes from the Royal Zoological Museum of the Netherlands at Leyden. 1879;1:165-84.
2. Subasinghe RP, Barg U, Tacon A. Chemicals in Asian aquaculture: need, usage, issues and challenges. In: Use of Chemicals in Aquaculture in Asia. Arthur JR, Lavilla-Pitogo CR, Subasinghe RP (eds). Southeast Asian Fisheries Development Center, Aquaculture Department Tigbauan, Iloilo, Philippines. 1996;1-6.
3. Gräslund S, Bengtsson BE. Chemicals and biological products used in South-East Asian shrimp farming and their potential impact on the environment: A review. *Sci Total Environ* 2001;280: 93-131.
4. Hopkins JS, Sandifer PA, Browdy CL. A review of water management regimes which abate the environmental impacts of shrimp farming. Proceedings of the Special Session on Shrimp Farming. Baton Rouge: World Aquaculture Society; 1995. p. 117-33
5. FDA. Food and Drug Administration. Chapter 11: Aquaculture Drugs in Fish and Fishery Products Hazards and Controls Guide. 2nded, Washington DC:FDA; 1998. p. 115-32.
6. FDA. New Animal Drugs Application. Freedom of Information Summary. Available from: <http://www.fda.gov/AnimalVeterinary/DevelopmentApprovalProcess/Aquaculture/ucm132954.htm> [Last accessed on 2007].
7. FDA. A quick reference guide to: Approved drugs for use in aquaculture. USFWS AADAP Program Available from: <http://www.fws.gov/fisheries/aadap>, the AFS Fish Culture Section Available from: <http://www.fishculturesection.org>. AFS Fish Health Section Available from: <http://www.fisheries.org/units/fhs/>. and FDA Center for Veterinary Medicine Available from: <http://www.fda.gov/cvm>. [Last accessed on 2008].
8. GEASAMP. Towards safe and effective use of chemicals in coastal aquaculture. GEASAMP Reports and Studies No. 65. Pub.IMO/ FAO/ UNESCO/ IOC/ WMO/ WHO/ IAEA/ UN/ UNEP, Rome:FAO; 1997
9. AftabUddin S, Abdul Kader Md. The use of antibiotics in shrimp hatcheries in Bangladesh. *J Fisheries Aquatic Sci* 2006;1:64-67.
10. Frappaolo PJ, Guest GB. Regulatory status of tetracyclines, penicillin and other antimicrobial microbial drugs in animal feeds. *Indian J Anim Sci* 1986;62:86-92.
11. Dierberg FE, Kiattisimkul W. Issues, impacts and implications of shrimp aquaculture in Thailand. *Environmental Management* 1996;20:649-66.
12. Bergheim A, Asgard T. Waste production from aquaculture. In: DJ Baird, MCCC Beveridge, LA Kelly, JF Muir, editors. *Aquaculture and Water Resource Management*. Institute of Aquaculture. Oxford, UK: Blackwell Science Ltd; 1996.

13. Alderman DJ, Rosenthal H, Smith P, Stewart J, Weston DP. Chemicals used in mariculture. ICES cooperative research report No. 202. Copenhagen, Denmark. ICES Working group. Environmental Interactions of Mariculture, 1994. p. 100.
14. Zitko V. Chemicals in aquaculture (an overview). *In: Proceedings of the Canada-Norway Workshop on Environmental Impacts of Aquaculture*, Institute of Marine Research. Fisker og havet NR 13, 1994.
15. Reardon IS, Harrell RM. Effects of varying salinities on the toxicity of potassium permanganate to larval and juvenile striped bass, *Morone saxatilis* (Walbaum). *Aquacult Fish Manage* 1994;25:571-8.
16. Curry S. Methemoglobinemia. *Ann Emerg Med* 1982;11:214-21.
17. Kirsch IR, Cohen HJ. Heinz body hemolytic anemia from the use of methylene blue in neonates. *J Pediatr* 1980;96:276-8.
18. Salaris SC, Babbs CF, Voorhees WD 3rd. Methylene blue as an inhibitor of superoxide generation by xanthine oxidase. *Biochem Pharmacol* 1991;42:499-506.
19. Katsuki S, Arnold W, Mittal C, Murad F. Stimulation of guanylatecyclase by sodium nitroprusside, nitroglycerin and nitric oxide in various tissue preparations and comparison to the effects of sodium azide and hydroxylamine. *J Cyclic Nucleotide Res* 1977;3:23-35.
20. Moncada S, Palmer RM, Higgs EA. Nitric oxide: Physiology, pathophysiology, pharmacology. *Pharmacol Rev* 1991;43:109-42.
21. Hampl V, Tristani-Firouzi M, Weir EK, Huang JM, Archer SL. Cyclic guanosine monophosphate as a hyperpolarizing factor. *In: Vanhoutte PM, editors. Endothelium-derived hyperpolarizing factor*. Amsterdam: Harwood Academic Publishers; 1996. p. 153-60.
22. Cragan JD. Teratogen Update: Methylene Blue. *Teratology* 1999;60:42-8.
23. Noga EJ. *Fish Disease, Diagnostic and Treatment*. Ames: Iowa State University Press; 1996.
24. Douet DG, Le Bris H, Giraud E. Environmental aspects of drug and chemical use in aquaculture: An overview. The use of veterinary drugs and vaccines in Mediterranean aquaculture, *Options Méditerranéennes*, 2009; A/ no. 86.

How to cite this article: Paul M, Chanda M, Maity J, Sen SS, Patra BC, Dash G. The use of antibiotics and disinfectants in the freshwater prawn (*Macrobrachium rosenbergii*) farms of north and south 24 Parganas districts of West Bengal. *Chron Young Sci* 2011;2:55-8.

Source of Support: Nil, **Conflict of Interest:** None declared

FORM IV

Statement about ownership and other particulars about newspaper (Chronicles of Young Scientists) to be published in the first issue every year after the last day of February

1. Place of publication : Mumbai
2. Periodicity of its publication : Quarterly (January, April, July and October)
3. Printer's Name : Medknow Publications & Media Pvt. Ltd.
Nationality : Indian
Address : B5-12, Kanara Business Center,
Off Link Rd, Ghatkopar (E), Mumbai - 400075, India.
Phone: 91-22-6649 1818
4. Publisher's Name : Dr. D. K. Sahu
For Medknow Publications & Media Pvt. Ltd.
Nationality : Indian
Address : B5-12, Kanara Business Center,
Off Link Rd, Ghatkopar (E), Mumbai - 400075, India.
Phone: 91-22-6649 1818
5. Editor's Name : Mr. Himanshu Gupta
Nationality : Indian
Address : Chronicles of Young Scientists
21, Jaina Building, Roshanara Road,
Delhi-110007, India
6. Names and addresses of individuals who own the newspaper and partners or shareholders holding More than one per cent of the total capital. : Organization of Pharmaceutical Unity with BioAllied Sciences (OPUBS)

I, **Mr. Himanshu Gupta** hereby declare that the particulars given above are true to the best of my knowledge and belief.

Date:

Dr. D. K. Sahu

Mr. Himanshu Gupta